

NOTES ON NEW AND NOTEWORTHY PLANTS. CV

Harold N. Moldenke

STACHYTARPHETA ANGUSTIFOLIA f. RIONEGRENSIS Moldenke, f. nov.

Haec forma a forma typica speciei serratura foliorum acutiora patentioreque et laminis perfragilibus recedit.

This form differs from the typical form of the species and its previously described New World forms in having its leaf-blades extremely thin and fragile, decidedly nigrescent in drying, the serration on the margins conspicuously more acute and wide-spreading, the individual teeth of varying size.

The type of the form was collected by Carl Friedrich Philipp von Martius in woods at M. Araraccara, along the Rio Negro, Roraima, Brazil, and is deposited in the herbarium of the Botanische Staatssammlung at Munich.

STACHYTARPHETA TRISPICATA var. OVATIFOLIA Moldenke, var. nov.

Haec varietas a forma typica speciei laminis foliorum latiore ovalibus ovatisve 4--4.5 cm. latis recedit.

This variety differs from the typical form of the species in having its leaf-blades distinctly more broadly oval or ovate, the mature ones 4--4.5 cm. wide at the widest point.

The type of the variety was collected by Edmundo Pereira (no. 9754; G. Pabst 8643) 41 km. from Vitória da Conquista, "rumo Anagé", Bahia, Brazil, on January 26, 1965, and is deposited in the herbarium of the Botanische Staatssammlung at Munich. The collector describes it as a shrub, 2--3 m. tall, with blue corollas.